1 NORMALIZING THE LEAST-SQUARES SOLUTION OF LINES

1.1 Description of the problem

Let a_1, b_1, c_1, k_1, a_2, b_2, c_2, k_2, a_3, b_3, c_3, k_3, be real numbers such that

$$a_1^2 + b_1^2 = a_2^2 + b_2^2 = a_3^2 + b_3^2 = 1.$$

We call d_1, d_2 and d_3 the straight lines of equations

$$\begin{cases} k_1a_1x_1 & + & k_1b_1x_2 & = & k_1c_1 \\ k_2a_2x_1 & + & k_2b_2x_2 & = & k_2c_2 \\ k_3a_3x_1 & + & k_3b_3x_2 & = & k_3c_3 \end{cases}$$

Note that in the Java Sketches the scaling numbers k_1, k_2 and k_3 are denoted by a, b and c.

NOTE FOR ERIC: IN JSXGRAPH FIGURES WE CAN USE SUBSCRIPTS!

The lines form a triangle of vertices $S_1 = d_2 \cap d_3$, $S_2 = d_3 \cap d_1$ and $S_3 = d_1 \cap d_2$, see Figure 1. We define



Figure 1: The triangle

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad \mathbf{c} = \begin{pmatrix} k_1c_1 \\ k_2c_2 \\ k_3c_3 \end{pmatrix}, \quad A = \begin{pmatrix} k_1a_1 & k_1b_1 \\ k_2a_2 & k_2b_2 \\ k_3a_3 & k_3b_3 \end{pmatrix}$$

The matrix equation $A\mathbf{x} = C$ has in general no solution. The least squares solution $\mathbf{x}_0$ of the normal equation

$$A^TA\mathbf{x} = A^T\mathbf{c}$$

is such that

$$F(\mathbf{x}) = F(x_1, x_2) = \|A\mathbf{x} - \mathbf{c}\|^2 = (\mathbf{x}^T\, A^T - \mathbf{c}^T)(A\mathbf{x} - \mathbf{c})$$

is minimum for $\mathbf{x} = \mathbf{x}_0$.

We denote by $d(\mathbf{x}, L)$ the distance from the point $\mathbf{x}$ to line L. If one computes $F(\mathbf{x})$ one gets:

$$F(\mathbf{x}) = k_1^2 \left(d(\mathbf{x}, d_1)\right)^2 + k_2^2 \left(d(\mathbf{x}, d_2)\right)^2 + k_3^2 \left(d(\mathbf{x}, d_3)\right)^2$$

I Case $k_1 = k_2 = k_3$

The function to be minimized is

$$F(\mathbf{x}) = \left(d(\mathbf{x}, d_1)\right)^2 + \left(d(\mathbf{x}, d_2)\right)^2 + \left(d(\mathbf{x}, d_3)\right)^2$$

Let us call $l_1 = S_2S_3$, $l_2 = S_3S_1$ and $l_3 = S_1S_2$, the lengths of the sides of the triangle defined by the lines d_1, d_2 and d_3. The point $\mathbf{x}_0$ such that $F(\mathbf{x})$ is minimum for $\mathbf{x} = \mathbf{x}_0$ is the barycenter of S_1, S_2 and S_3 with masses l_1^2, l_2^2 and l_3^2:

$$\mathbf{x}_0 = \frac{l_1^2}{l_1^2 + l_2^2 + l_3^2} S_1 + \frac{l_2^2}{l_1^2 + l_2^2 + l_3^2} S_2 + \frac{l_3^2}{l_1^2 + l_2^2 + l_3^2} S_3$$

Problems

1. If the triangle defined by the lines d_1, d_2 and d_3 is isosceles (let us say $S_1S_2 = S_1S_3$), on what line do you expect to find $\mathbf{x}_0$?



Figure 2: Isosceles triangle

2. If the triangle is equilateral where is $\mathbf{x}_0$?

3. The distances of $\mathbf{x}_0$ to the sides of the triangle are proportional to the lengths of the corresponding sides to a certain power r:

$$\frac{d(\mathbf{x}_0, d_1)}{{l_1}^r} = \frac{d(\mathbf{x}_0, d_2)}{{l_2}^r} = \frac{d(\mathbf{x}_0, d_3)}{{l_3}^r}$$

Guess what is the value of r?

Note: If $r = -1$, the point $\mathbf{x}_0$ would be the center of gravity; if $r = 0$, the point $\mathbf{x}_0$ would be the center of the inner circle.

4. If the triangle is rectangular, e.g. $d_2 \perp d_3$, on what line do you find $\mathbf{x}_0$ (see Figure 3?



Figure 3: Rectangular triangle

$$\frac{HK_2}{S_1S_3} = \frac{HK_3}{S_2S_1} \text{ since the triangles are similar}$$

II Case k_1, k_2 k_3 any real numbers

For this general case we already made experiments!

NOTE FOR ERIC: ONE VERSION OF THE DOCUMENT ENDED HERE

1. Does the triangle $S_1S_2S_3$ change if you change the k_i's ?

2. How does $\hat{\mathbf{x}}$ move if you change k_1, keeping all other numbers fixed?

3. Where is $\hat{\mathbf{x}}$ if the triangle is isosceles ($l_2 = l_3$) and $k_2 = k_3$?

4. Where is $\hat{\mathbf{x}}$ if $k_1 = 0$?

5. To a point of what side of the triangle does $\hat{\mathbf{x}}$ tend to, when $k_1 \to \infty$, keeping all other numbers fixed?

Is it possible to choose k_2 and k_3 in such a way that $\hat{\mathbf{x}}$ tends to the middle of one side?

Answer: "Yes, $k_2 = 1/l_2$, $k_3 = 1/l_3$? I'm not sure (??)."

6. If you use a scaling matrix

$$D = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$$

to compute the solution of the scaled least square equation

$$A^T D^T D A\mathbf{x} = A^T D^T D\mathbf{c}$$

How should you choose D to get the solution $\hat{\mathbf{x}}$ of the case I?

Answer: "$\lambda_i = 1/k_i$."

Never forget: always scale before using least squares solution to avoid impact of the choice of units.

1.2 A "geometric" construction of $\hat{\mathrm{x}}$ in the case I



1. Construct B_1 such that $B_1C = CA$ and $B, C \perp CA$ and B_1 and B on the same side of AC:



2 Construct the line through B_1, parallel to AC, call it e.

3 Construct C_1 such that $BC_1 = BA$, $BC_1 \perp BA$, C_1 and C on the same side of AB.

4 Construct the line f through C_1, parallel to AB.

5 Let O be the intersection of e and f.

6 Join AO. The point $\hat{\mathrm{x}}$ has to be on AO.





7 Do the same construction from the vertex B, getting BO'.

8 $\hat{\mathrm{x}}$ is the intersection of AO and BO'.

9 Verify with a construction CO'' through C.
The points C, $\hat{\mathrm{x}}$ and O'' should be on a straight line.


B''
O''
O'
A
B
C
A''
O